

Tensor Product of Vertex Operator Algebras

Antun Milas

17.8.1995

Abstract

Let $V=V_1 \otimes V_2$ be a tensor product of VOAs. Using Zhu theory we discuss the theory of representations of V (associative algebra, modules and 'fusion rules'). We prove that this theory is more or less the same as representation theory of tensor product of the associative algebras.

1 Introduction

In this paper we discuss some things which are closely related with so called aksiomatic approach to VOA [FHL], [FLM], [B], [Li] [DL]. There are lot of literature in theory of VOAs, representation theory etc. It seems that Borchards [B] first make a structure of VOA on tensor product on very natural way. It's not worth to mention that construction of tensor product of modules is rather complicated [Li] and use the theory of intertwining operators.

In this paper we gave some general results about tensor product of VOAs. We find that associative algebra ([Z], [FZ]) have a good functorial property related to tensor product of two VOAs (Theorem 4.1).

The other aspect which is considered is representation theory (classification of modules and calculating the 'fusion rules'). Using the upper property

*1991 Mathematics Subject Classification. Primary 17b65; Secondary 17A70, 17B67.

of associative algebra we prove the classification theorem (see [FHL]) and discuss whether tensor product of two VOAs is a rational VOA.

Using the structure of bimodules (section 5.) we present the proof of fusion rules formula for $V = V_1 \otimes V_2$.

It's easy to make the similar theory in case of tensor product of two super-algebras [KWn], [Li1]. Also the similar results are valid in the classification theory of twisted modules of tensor product of VOA [DLM].

The next step closely related with this theory would be consideration of *dual pairs* in VOAs (see [FZ]), or *coset* models (WZNW models, minimal models). If we have VOA V and V_1 and V_2 subalgebras which 'commute' then we said that (V_1, V_2) form a dual pair for V -module M if $M = \oplus_i M_i \otimes N_i$, where $M_i(N_i)$ are V_1 (V_2)-modules. We are working on this problematic.

2 Preliminaries

Vertex operator algebras is mathematical counterpart of Conformal field theory (CFT). Today there are various definitions of VOA and representations of VOA according to graduation, Virasoro element etc..So we're going to define VOA and modules in generality what we need.

2.1 Vertex operator algebras and modules

Definition 2.1 *A vertex operator algebra is a $\mathbf{Z}$ -graded vector space $V = \bigoplus_{n \in \mathbf{Z}} V_n$ with a sequence of linear operators $\{a(n) \mid n \in \mathbf{Z}\} \subset \text{End } V$ associated to every $a \in V$, such that for fixed $a, b \in V$, $a(n)b = 0$ for n sufficiently large. We call the generating series $Y(a, z) = \sum_{n \in \mathbf{Z}} a(n)z^{-n-1} \in (\text{End } V)[[z, z^{-1}]]$, vertex operators associated to a , satisfy the following axioms:*

(V1) $Y(a, z) = 0$ iff $a = 0$.

(V2) *There is a vacuum vector, which we denote by $\mathbf{1}$, such that*

$$Y(\mathbf{1}, z) = I_V \text{ (} I_V \text{ is the identity of } \text{End } V \text{)}.$$

(V3) There is a special element $\omega \in V$ (called the Virasoro element), whose vertex operator we write in the form

$$Y(\omega, z) = \sum_{n \in \mathbf{Z}} \omega(n) z^{-n-1} = \sum_{n \in \mathbf{Z}} L_n z^{-n-2},$$

such that

$$L_0 |_{V_n} = nI |_{V_n},$$

$$Y(L_{-1}a, z) = \frac{d}{dz} Y(a, z) \quad \text{for every } a \in V, \quad (1)$$

$$[L_m, L_n] = (m - n)L_{m+n} + \delta_{m+n,0} \frac{m^3 - m}{12} c, \quad (2)$$

where c is some constant in $\mathbf{C}$, which is called the rank of V .

(V4) The Jacobi identity holds, i.e.

$$\begin{aligned} & z_0^{-1} \delta \left(\frac{z_1 - z_2}{z_0} \right) Y(a, z_1) Y(b, z_2) - z_0^{-1} \delta \left(\frac{-z_2 + z_1}{z_0} \right) Y(b, z_2) Y(a, z_1) \\ &= z_2^{-1} \delta \left(\frac{z_1 - z_0}{z_2} \right) Y(Y(a, z_0)b, z_2) \end{aligned} \quad (3)$$

for any $a, b \in V$.

Definition 2.2 Let V_1 and V_2 be two VOAs then mapping $f : V_1 \rightarrow V_2$ is homomorphism of VOAs if F is a homomorphism of vector spaces and if $f(Y(a, z)b) = Y(f(a), z)f(b)$ for every $a, b \in V$. f is an isomorphism of VOAs if f is bijection.

Let V_1 and V_2 be two VOAs. Then the space $V_1 \oplus V_2$ has a natural VOA structure. The subspace I of V is called ideal if $Y(a, z)b \in I[[z, z^{-1}]]$ for every $a \in V, b \in I$. Given an ideal I in V such that $\mathbf{1} \notin I$, $\omega \notin I$, the quotient V/I admits a natural VOA structure (see [FZ]). VOA is simple if it doesn't have nontrivial ideals. Also, we define semisimple VOA as VOA which is direct sum of simple VOAs.

Definition 2.3 Given an VOA V , a representation of V (or V-module) is a $\mathbf{Z}_+$ -graded vector space $M = \bigoplus_{n \in \mathbf{Z}_+} M_n$ and a linear map

$$V \longrightarrow (\text{End } M)[[z, z^{-1}]],$$

$$a \longmapsto Y_M(a, z) = \sum_{n \in \mathbf{Z}} a(n) z^{-n-1},$$

satisfying

(M1) $a(n)M_m \subset M_{m+\deg a-n-1}$ for every homogeneous element a .

(M2) $Y_M(1, z) = I_M$, and setting $Y_M(\omega, z) = \sum_{n \in \mathbf{Z}} L_n z^{-n-2}$, we have

$$[L_m, L_n] = (m-n)L_{m+n} + \delta_{m+n,0} \frac{m^3-m}{12} c, \text{ nonnumber} \quad (4)$$

$$Y_M(L_{-1}a, z) = \frac{d}{dz} Y_M(a, z)$$

for every $a \in V$.

(M3) The Jacobi identity holds, i.e.

$$z_0^{-1} \delta \left(\frac{z_1 - z_2}{z_0} \right) Y_M(a, z_1) Y_M(b, z_2) - z_0^{-1} \delta \left(\frac{-z_2 + z_1}{z_0} \right) Y_M(b, z_2) Y_M(a, z_1) \\ z_2^{-1} \delta \left(\frac{z_1 - z_0}{z_2} \right) Y_M(Y(a, z_0)b, z_2) \quad (5)$$

for any $a, b \in V$.

Remark 2.1 In [FHL] the definition of V -module is stronger, with the following additional aksiom:

(M0) Dimension of $M_0(n)$ (graded subspace) is finite for every n .

2.2 Intertwinig operators

Definition 2.4 Let M^1, M^2 and M^3 be three V -modules. An intertwinig operator of type $\begin{pmatrix} M^3 \\ M^1, M^2 \end{pmatrix}$ is a linear map

$$I(, x) : M^1 \rightarrow (Hom(M^2, M^3)\{x\}) \quad u \mapsto I(u, z) = \sum_{\alpha \in \mathbf{C}} u_{\alpha} z^{-\alpha-1} \quad (6)$$

satisfying the following conditions:

- (I1) For any fixed $u \in M^1$, $v \in M^2$, $\alpha \in C$, $u_{\alpha+n}v = 0$ for n sufficiently large;
- (I2) $I(L(-1)u, z)v = \frac{d}{dz}I(u, z)v$ for $u \in M^1$, $v \in M^2$;
- (I3) The Jacobi identity holds, i.e.

$$\begin{aligned} & z_0^{-1} \delta \left(\frac{z_1 - z_2}{z_0} \right) Y_{M^3}(a, z_1) I(u, z_2)v - z_0^{-1} \delta \left(\frac{-z_2 + z_1}{z_0} \right) I(u, z_2) Y_{M^2}(a, z_1)v = \\ & z_2^{-1} \delta \left(\frac{z_1 - z_0}{z_2} \right) I(Y(a, z_0)u, z_2)v \end{aligned} \quad (7)$$

for any $a \in V$, $u \in M^1$, $v \in M^2$.

Denote by $I \begin{pmatrix} M^3 \\ M^1, M^2 \end{pmatrix}$ the vector space of all intertwining operators of this type. Dimension of this space we call *fusion rules* of corresponding type.

By using the definition of fusion rules we can define (under some circumstances) so called *fusion algebra* [DL], [Z]. It's very interesting that in many examples like WZNW models ([DL], [FZ], [Li1]) minimal models ([W], [KWn]) the fusion rules are the same as in CFT definition of 'fusion' and also in geometrical definition .

3 Tensor product

Let V_1 and V_2 be two vertex operator algebras. Following [B] and [FHL] we introduce the VOA structure on vector space $V=V_1 \otimes V_2$.

Firstly, $V_1 \otimes V_2$ is on natural way $\mathbf{Z}_{\geq 0}$ -graded vector space with natural graduation. We define the field associated to $a \otimes b \in V$ as follows: $Y(a \otimes b, z) = Y(a, z) \otimes Y(b, z)$. If ω_1 and ω_2 are the Virasoro elements of V_1 and V_2 it's easy to see that vector $\omega_1 \otimes \mathbf{1} + \mathbf{1} \otimes \omega_2$ is a Virasoro element for V .

The axioms [V1], [V2] and [V3] are satisfied so (with this definition) V is a VOA.

Now the following simple lemma holds

Lemma 3.1 *Let V_1 V_2 and V_3 be a three VOAs. Then :*

$$(1) \ V_1 \otimes (V_2 \otimes V_3) \cong (V_1 \otimes V_2) \otimes V_3$$

$$(2) \ V_1 \otimes V_2 \cong V_2 \otimes V_1$$

Also we need this fact.

Proposition 3.1 *[FHL] Let M_1 (M_2) be irreducible V_1 (V_2)-module with $M(0)$ property. Then $M_1 \otimes M_2$ is irreducible $V_1 \otimes V_2$ -module.*

4 The associative algebra $A(V)$

4.1 Definition and properties

Let V be a VOA. For any homogeneous element $a \in V$ and for any $b \in V$, following [Z] we define

$$a * b = \text{Res}_z \frac{(1+z)^{\text{wta}}}{z} Y(a, z)b. \quad (8)$$

$$a \circ b = \text{Res}_z \frac{(1+z)^{\text{wta}}}{z^2} Y(a, z)b. \quad (9)$$

Then extend this products bilinearly to the whole space V . Let $O(V)$ be a subspace of V linearly spanned with set $\{a \circ b : a, b \in V\}$.

Set $A(V) = V/O(V)$. The multiplication $*$ induces the multiplication on the $A(V)$ and $A(V)$ becomes an associative algebra. The image of $\mathbf{1}$ in $A(V)$ becomes the identity element till the image of ω is in center of $A(V)$ (see [Z]).

4.2 Structure of $A(V_1 \otimes V_2)$

As we saw in previous section to every VOA we associate the associative algebra $A(V)$. Here we are going to prove that associative algebras $A(V_1 \otimes V_2)$ and $A(V_1) \otimes A(V_2)$ are isomorphic.

First we prove some auxiliary results:

Proposition 4.1 *Let $a, b \in V$ then*

$$a \circ_m b := \text{Res}_z \frac{(1+z)^{\text{wta}}}{z^m} Y(a, z)b \in O(V),$$

for every $m \geq 2$.

Proof. Put $a' = L(-1)^{m-1}a$. Then it's easy to see that $a' \circ b = C(a \circ_m b)$ for some constant C .

Lemma 4.1 *Let $a, a_1 \in V_1$ and $b \in V_2$ then*

$$(a \otimes \mathbf{1}) \circ (a_1 \otimes b) = (a \circ a_1) \otimes b$$

Proof. From the fact $\mathbf{1}_n a = \delta_{-1,n} a$ and from the definition of bilinear mapping $\circ$ it follows:

$$(a \otimes \mathbf{1}) \circ (a_1 \otimes b) = \sum_{m \in \mathbb{Z}} (a \circ_m a_1) \otimes \mathbf{1}_{-m+1} b = (a \circ a_1) \otimes b.$$

Lemma 4.2

$$O(V_1 \otimes V_2) = O(V_1) \otimes V_2 + V_1 \otimes O(V_2). \quad (10)$$

Proof. From Lemma 4.1 follows that $O(V_1) \otimes V_2 + V_1 \otimes O(V_2) \subseteq O(V_1 \otimes V_2)$. The reverse direction follows from the definition of $\circ_m$ and Proposition 4.1.

Theorem 4.1 *We define the mapping*

$$F : A(V_1 \otimes V_2) \rightarrow A(V_1) \otimes A(V_2), \quad (11)$$

$$[a \otimes b] \mapsto [a] \otimes [b]. \quad (12)$$

The mapping F is isomorphism of associative algebras with identity.

Proof. Because of

$$A(V_1) \otimes A(V_2) = (V_1 \otimes V_2)/(O(V_1) \otimes V_2 + V_1 \otimes O(V_2)) \quad (13)$$

and (10) it follows that this mapping is well-defined. This map is also surjective.

We want to prove injectivity. Let $m = \sum_i [a_i \otimes b_i]$ and $F(m) = 0$, then $\sum_i a_i \otimes b_i \in O(V_1) \otimes V_2 + V_1 \otimes O(V_2)$, and from (13) $\sum_i a_i \otimes b_i \in O(V_1 \otimes V_2)$ follows that $\sum_i [a_i \otimes b_i] = 0$, so we prove injectivity.

We denote the multiplication in $A(V_1) \otimes A(V_2)$ with the same symbol $*$. Now,

$$\begin{aligned} F([a \otimes b] * [a' \otimes b']) &= \text{Res}_z[Y(a \otimes b, z) \frac{(1+z)^{\text{wta}+\text{wtb}}}{z} a' \otimes b'] \\ &= [\text{Res}_z(Y(a, z) \frac{(1+z)^{\text{wta}}}{z} a' \otimes Y(b, z)(1+z)^{\text{wtb}} b')] \\ &= \sum_{i \in \mathbf{Z}} [a \circ_{-i} a' \otimes b \circ_{i+2} b'] \\ &= (\text{by Proposition 4.1}) ([a] * [a']) \otimes ([b] * [b']) \\ &= ([a] \otimes [b]) * ([a'] \otimes [b']) \\ &= (\text{by Lemma 4.2}) F([a \otimes b]) * F([a' \otimes b']). \end{aligned} \quad (14)$$

so F is homomorphism.

5 Modules and fusion rules for $V_1 \otimes V_2$

Among all VOAs the most interesting are 'rational VOAs'.

Definition 5.1 *[Z], [FZ] VOA is rational if it have only finitely many irreducible representations and if every finitely generated module for V is completely reducible. For the VOA which have only finitely many irreducible modules we said that is quasirational.*

5.1 Zhu's theorem

Let $M = \oplus_{n \in \mathbf{Z}_+} M_n$ be a V -module. For a homogeneous element $a \in V$ we define $o(a) = a(\deg a - 1)$. From the definition of M follows that operator $o(a)$ preserves the gradation of M .

Theorem 5.1 [FZ]

(a) On $\text{End}(M_0)$ we have

$$o(a)o(b) = o(a * b)$$

$$o(x) = 0$$

for every $a, b \in V$, $x \in O(V)$. The top level M_0 is an $A(V)$ -module.

(b) Let U be an $A(V)$ -module, there exists V -module M such that $A(V)$ -module M_0 and U are isomorphic.

Thus, we have one-to-one correspondence between irreducible V -modules and irreducible $A(V)$ -modules.

Remark 5.1 There is a explicit way how to every $A(V)$ -module we can associate V -module. Moreover, if $L(0)$ acts semisimple then for semisimple $A(V)$ -module M corresponding V -module is semisimple ([Li],[Z],[DLM]).

5.2 Classification theorem

Put $V = V_1 \otimes V_2$. In this paragraph we give a the proof of Frenkel-Huang-Lepowsky theorem of classification of all irreducible V -modules with some weaker condition by using different approach ($A(V)$ -theory).

Remark 5.2 We will (from now till the end) assume that $L(0)$ acts semisimply on every V -module.

Now we need prove technical lemma about representations of tensor product of two associative algebras with identity. The proof is from [W].

Lemma 5.1 Let A and B be two associative algebras with identity and M a finite-dimensional irreducible $A \otimes B$ -module. Then $M \cong M_1 \otimes M_2$ where M_1 (M_2) is irreducible module for A (B).

Proof. Let M_1 be an irreducible submodule of V (for A). Then we are looking the sum $\sum_{b \in B} bM_1$ (b means $1 \otimes b$). This sum is a whole space M because M is irreducible $A \otimes B$ -module. Now M is (as A -module) completely reducible and every irreducible submodule is isomorphic to M_1 (this follows from Schur's Lemma). Set $M_2 = \text{Hom}_A(M_1, M)$. Then M_2 is on natural way B -module. The module $M' = M_1 \otimes M_2$ is $A \otimes B$ -module. Now we have $A \otimes B$ -homomorphism from M' to a submodule of M such that

$$a \otimes f \mapsto f(a).$$

This homomorphism is bijection so it's isomorphic to M .

Remark 5.3 *From Burnside's theorem follows that every $A \otimes B$ -module of the form $M_1 \otimes M_2$ is irreducible if M_1 and M_2 are finite-dimensional. If M_1 or M_2 is infinite-dimensional Burnside's theorem doesn't implies the irreducibility.*

The following corollary is a modification of [FHL].

Corrolary 5.1

- (1.) *If V_1 and V_2 are two VOA then every irreducible V -module M with $M(0)$ property is isomorphic to $M_1 \otimes M_2$ where M_1 and M_2 are irreducible V_1 (V_2)-modules.*
- (2.) *If $A(V_1)$ and $A(V_2)$ are finite-dimensional then every irreducible V -module is of the form $M_1 \otimes M_2$ where M_1 and M_2 are irreducible.*

Proof. Follows directly from Theorem 4.1, Theorem 5.1, Lemma 5.1 and Lemma 3.1.

Remark 5.4 *In some cases of VOA (WZNW models [FZ],[KWn], minimal models [W]) associative algebra $A(V)$ is a finite dimensional (and rational).*

From the definition of rationality follows:

Lemma 5.2 *If V is a rational VOA then $A(V)$ is finite-dimensional semisimple associative algebra with identity.*

This fact is also proved in [DLM] with weaker definition of rationality.

Now we are going to discuss rationality of V if V_1 and V_2 are rational. From Remark 5.1 we must analyze when finitely generated $A(V)$ -module is semisimple. This is equivalent to the fact that every $A(V)$ -module is semisimple. Similar problem was discussed in [DLM1].

Theorem 5.2 *If V_1 and V_2 are two rational VOAs then $V = V_1 \otimes V_2$ is also rational. If V_1 is rational and $A(V_2)$ is finite dimensional then $V = V_1 \otimes V_2$ is quasirational.*

Proof. Second fact follows directly from the Corrolary 5.1. If V_1 and V_2 are rational V is from the second fact quasirational VOA. But tensor product of two finite dimensional semisimple algebras is also semisimple algebra so every finitely generated module is semisimple ([P],[J]) and V is rational.

5.3 Fusion rules

Let $V_1 \otimes V_2$ be VOA and $M^i = M_1^i \otimes M_2^i$, $i = 1, 2, 3$ its three irreducible modules such that $L_1(0)$, $L_2(0)$ act semisimply on M_1^i , M_2^i $i=1,2,3$.

First, we need the following fact

Proposition 5.1 *[Li],[DL] Let $I(\cdot, z)$ be an intertwining operator of type $\begin{pmatrix} M^3 \\ M^1, M^2 \end{pmatrix}$ then the Jacobi identity is equivalent to following conditions :*
For every $a \in V$, $v \in M_1$, $w \in M_2$ exists $r, k \in \mathbf{N}$ such that ('associativity')

$$(z_0 + z_2)^r Y(a, z_0) I(v, z_2) w = (z_0 + z_2)^r I(Y(a, z_0) v, z_2) w$$

and ('commutativity')

$$(z_0 - z_1)^k Y(a, z_0) I(v, z_2) w = (z_0 - z_1)^k I(v, z_2) Y(a, z_0) w$$

are satisfied.

Now using this description of 'Jacobi identity' for intertwinig operators we have the following theorem.

Proposition 5.2 *Let V and M^i , $i = 1, 2, 3$ be as above then*

$$\dim I \begin{pmatrix} M^3 \\ M^1, M^2 \end{pmatrix} \geq \dim I \begin{pmatrix} M_1^3 \\ M_1^1, M_1^2 \end{pmatrix} \dim I \begin{pmatrix} M_2^3 \\ M_2^1, M_2^2 \end{pmatrix}.$$

Proof. If I_1 is an intertwining operator of type $\begin{pmatrix} M_1^3 \\ M_1^1, M_1^2 \end{pmatrix}$ and I_2 of type $\begin{pmatrix} M_2^3 \\ M_2^1, M_2^2 \end{pmatrix}$ then we define $I(, z)$ as on the following way:

$$I(m_1 \otimes m_2, z)n_1 \otimes n_2 = I_1(m_1, z)n_1 \otimes I_2(m_2, z)n_2, \quad (15)$$

for every $m_1 \otimes m_2 \in M^1$ and $n_1 \otimes n_2 \in M^2$. Now the first two property ([I1] and [I2]) of intertwining operators are obviously satisfied. To prove Jacobi identity we use lemma 5.1. Associativity and commutativity we prove on following way.

If associativity for I_1 is satisfied for some $r_1 \in \mathbf{N}$, $a_1 \in V_1$, $v_1 \in M_1$ and I_2 for some $r_2 \in \mathbf{N}$, $a_2 \in V_2$, $v_2 \in M_2$. Then for $u = a_1 \otimes a_2$ and $v = v_1 \otimes v_2$ we put $r = \max\{r_1, r_2\}$ and associativity is satisfied. In similar way we can prove commutativity. Now it is easy to see that proposition is true.

The equality in Proposition 5.2 is not satisfied in general but in some cases it will hold.

5.4 The structure of bimodules

Let $M = M_1 \otimes M_2$ be a $V = V_1 \otimes V_2$ -module. Then we define $A(M) = M/O(M)$ where $O(M) = \{a \circ m : a \in V, m \in M\}$. We define left and right action of $A(V)$ on $A(M)$ (see [Z]):

$$a \cdot m = [Res_z \frac{(1+z)^{wta}}{z} Y(a, z)m],$$

$$m \cdot a = [Res_z \frac{(1+z)^{wta-1}}{z} Y(a, z)m]$$

for every $a \in A(V)$, $m \in A(M)$. With this notation $A(M)$ is a $A(V)$ -bimodule.

Theorem 5.3 *Let M_1 (M_2) be irreducible V_1 (V_2)-modules. Let $a \in A(V_1)$, $b \in A(V_2)$ and $m_1 \in A(M_1)$, $m_2 \in A(M_2)$ then*

(1)

$$(a \otimes b) \cdot (m_1 \otimes m_2) = (a \cdot m_1) \otimes (b \cdot m_2),$$

(2)

$$(m_1 \otimes m_2) \cdot (a \otimes b) = (m_1 \cdot a) \otimes (b \cdot m_2).$$

Moreover relations (1) and (2) define an isomorphism of $A(M_1 \otimes M_2)$ and $A(M_1) \otimes A(M_2)$ as $A(V_1) \otimes A(V_2)$ -bimodules.

Proof.

From the irreducibility of M_1 and M_2 follows that $Y_{M_1}(\mathbf{1}, z)$ acts as a scalar on M_1 (same thing for M_2). Then it's not hard to prove:

$$O(M_1 \otimes M_2) = O(M_1) \otimes M_2 + M_1 \otimes O(M_2),$$

using the proofs of Lemmas 4.1 and 4.2.

The proof of (1) is analogous as the proof of Theorem 4.1. We want to prove (2) which with upper relation would give us the result.

We have

$$\begin{aligned} & (m_1 \otimes m_2) \cdot (a \otimes b) \\ &= Res_z \left(\frac{(1+z)^{wta+wtb-1}}{z} Y(a, z)m_1 \otimes Y(b, z)m_2 \right) \\ &= \sum_{i \in \mathbf{Z}} (Res_z(1+z)^{wta-1} z^{-i} Y(a, z)m_1 \otimes Res_z(1+z)^{wtb} z^{i-2} Y(b, z)m_2) \\ &= \sum_{i > 0} (Res_z(1+z)^{wta-1} z^{-i} Y(a, z)m_1 \otimes Res_z(1+z)^{wtb} z^{i-2} Y(b, z)m_2) \mod O(M_1 \otimes M_2) \\ &= Res_z(1+z)^{wta-1} z^{-1} Y(a, z)m_1 \otimes Res_z(1+z)^{wtb} z^{-1} Y(b, z)m_2 \\ &+ \sum_{i > 1} (Res_z(1+z)^{wta-1} (z^{-i} + z^{-i+1} - z^{-i+1}) Y(a, z)m_1 \\ &\otimes Res_z(1+z)^{wtb} z^{i-2} Y(b, z)m_2) \mod O(M_1 \otimes M_2) \\ &= m_1 \cdot a \otimes Res_z(1+z)^{wtb} z^{-1} Y(b, z)m_2 \\ &+ \sum_{i > 1} (Res_z((1+z)^{wta} z^{-i} - (1+z)^{wta-1} z^{-i+1}) Y(a, z)m_1 \\ &\otimes Res_z(1+z)^{wtb} z^{i-2} Y(b, z)m_2) \mod O(M_1 \otimes M_2) \\ &= m_1 \cdot a \otimes Res_z(1+z)^{wtb} z^{-1} Y(b, z)m_2 - Res_z(1+z)^{wta-1} z^{-1} Y(a, z)m_1 \\ &\otimes Res_z(1+z)^{wtb} Y(b, z)m_2 - \sum_{i > 2} (1+z)^{wta-1} z^{-i+1} Y(a, z)m_1 \end{aligned}$$

$$\begin{aligned}
& \otimes \text{Res}_z(1+z)^{wtb} z^{i-2} Y(b, z) m_2 \bmod O(M_1 \otimes M_2) \\
&= m_1 \cdot a \otimes \sum_{i>0} \text{Res}_z(1+z)^{wtb} z^{i-2} Y(b, z) m_2 \bmod O(M_1 \otimes M_2) \\
&= m_1 \cdot a \otimes \text{Res}_z(1+z)^{wtb} \frac{1}{z(1+z)} Y(b, z) m_2 \bmod O(M_1 \otimes M_2) \\
&= (m_1 \cdot a) \otimes (m_2 \cdot b) \bmod O(M_1 \otimes M_2) \\
&= (m_1 \cdot a) \otimes (m_2 \cdot b).
\end{aligned}$$

(16)

In the summation we use the fact that for $n \gg 0$

$$\text{Res}_z(1+z)^{wta} z^n Y(a, z) m_1 = 0,$$

so we have the proof.

We need the following formula for calculating 'fusion rules'.

Theorem 5.4 [FZ] *Let M_1, M_2 and M_3 be three V -modules such that $L(0)$ acts semisimple. There is one to one correspondance between*

$$\text{Hom}_{A(V)}(A(M_1) \otimes_{A(V)} M_2(0), M_3(0)) \quad (17)$$

and 'fusion rules' of corresponding type $(I \begin{pmatrix} M_3 \\ M_1, M_2 \end{pmatrix})$.

Lemma 5.3 *Let V be a rational VOA M_1, M_2 and M_3 its irreducible such that every 'fusion rules' is finite-dimensional then vector space*

$$M = A(M_1) \otimes_{A(V)} M_2(0) \quad (18)$$

is also finite-dimensional.

Proof. From the rationality of V follows that M is semisimple and then can be written:

$$M = \oplus_{i \in I} M_i, \quad (19)$$

where M_i are isotypical components $A(V)$ -module M . But $A(V)$ has only finite number of inequivalent irreducible modules so set I is finite. If some M_i is infinite-dimensional, and N term from M_i then $\text{Hom}_{A(V)}(M, N)$ would be infinite-dimensional which contradict with finiteness of 'fusion rules'.

Now we need following fact from tensor algebra.

Lemma 5.4 *Let A, B be two associative algebras with identity over $\mathbf{C}$ and M and N finite-dimensional right $A, (B)$ -modules and $M', (N')$ finite dimensional left $A, (B)$ -modules then:*

$$\begin{aligned} & \dim(M \otimes_A M') \otimes (N \otimes_B N') \\ &= \dim(M \otimes M') \otimes_{A \otimes B} (N \otimes N'). \end{aligned} \quad (20)$$

Using this Lemma we have:

Theorem 5.5 *(Same notation like in Proposition 5.2 If V_1 and V_2 are rational VOAs with finite 'fusion rules' then in Proposition 5.2 stands equality.*

Proof.

From [KWn] we have:

$$\begin{aligned} & \dim \text{Hom}_{A(V)}(A(M_1) \otimes_{A(V)} M_2(0), M_3(0)) \\ &= \dim M_3(0)^* \otimes_{A(V)} A(M_1) \otimes_{A(V)} M_2(0). \end{aligned} \quad (21)$$

Now if we put $A(V) = A(V_1) \otimes A(V_2)$, $M_2(0) = M_1^1(0) \otimes M_2^2(0)$, $M_3(0) = M_3^1(0) \otimes M_3^2(0)$ and $A(M_1) = A(M_1^1 \otimes M_1^2)$ we have from Lemma 5.4, 5.3 and Theorem 5.3

$$\begin{aligned} & \dim M_3(0)^* \otimes_{A(V)} A(M_1) \otimes_{A(V)} M_2(0) \\ &= \dim M_3^1(0)^* \otimes_{A(V_1)} A(M_1^1) \otimes_{A(V_1)} M_2^1(0) \\ & \quad \dim M_3^2(0)^* \otimes_{A(V_2)} A(M_1^2) \otimes_{A(V_2)} M_2^2(0). \end{aligned} \quad (22)$$

So we have the proof.

Corrolary 5.2 *If we have two rational VOAs with Zhu conditions and with 'fusion rules' 0 or 1 then tensor product have also 'fusion rules' 0 or 1.*

References

- [B] R.E. Borcherds, Vertex algebras, Kac-Moody algebras and the monster, Proc.Nat.Acad.Sci.USA, 83 (1986), 3068-3071
- [DL] C.Dong, J.Lepowsky, Generalized Vertex algebras and relative vertex operators, Progress in Math. Vol. **112**, Birkhauser, Boston 1993
- [DLM] C.Dong, H.Li, G.Mason, Twisted representations of VOAs, preprint
- [DLM1] C.Dong, H.Li, G.Mason, Regularity of rational VOAs, preprint
- [FHL] I.Frenkel, Y.-Z.Huang, J.Lepowsky, On axiomatic approach to Vertex operator algebras and modules, Memoirs of the AMS, **494**, 1994
- [FLM] I.Frenkel, J.Lepowsky, A.Meurman, Vertex operator algebras and the monster, Pure and Appl. Math.Vol. **134**, Academic Press, Boston, 1988
- [FZ] I.Frenkel, Y.-C.Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, Duke Math.J.**66**, 1992, 123-168
- [J] N.Jacobson, Strojenije koljca, Mir, Moscow 1964
- [K] V.Kac, Infinite dimensional Lie algebras, Cambridge Univ. Press, 1990
- [KWn] V.Kac, W.Wang, Vertex operator superalgebras and their representations, preprint
- [Li] H.Li, Representation theory and tensor product theory for Vertex Operator Algebras, PhD. thesis, Rutgers, 1994
- [Li1] H.Li, Local system of vertex operator algebras and superalgebras, preprint 1994
- [MP] A.Meurman, M.Primec, Annihilating fields of $sl(2, \mathbf{C})^\sim$ and combinatorial identities, preprint
- [P] R.Pierce, Associativnjie algeabri, Mir, Moscow 1986
- [W] W.Wang, Rationality of Virasoro Vertex operator algebra, Duke Math.J. IMRN, Vol **71**, No.1 (1993), 197-211
- [Z] Y.-C.Zhu, Vertex operator algebras, elliptic function and modular forms, PhD Thesis, Yale University, 1990.

Department of Mathematics, University of Zagreb, Bijenička 30, 41000 Zagreb, Croatia
e-mail : milas@cromath.math.hr